

2005

- I/1 : (a) $1/3$ (b) $1/6$ (c) -1
 2. (a) true (b) false (c) true (d) true (e) true but how to prove w/ suffix notation?
3. (a) $(1-p+np)^{-1}$ (b) (i) $1/365$ (ii) $N=254$ (c) 1.46
 4. (a) (i) $1-e^{-xt_0}$ (ii) Mean = $1/\alpha$ Var = $1/\alpha^2$ (iii) e^{-1} (b) (ii) $\sigma = 2C/\alpha$
5. (a) $\cos^{-1} 4/5$ (b) $\cos^{-1}(3/\sqrt{2})$ (c) $\sqrt{18}/2$ (d) $3/\sqrt{2}$
6. (a) - (b) and (c) $8\pi a^3$
 7. (a) - (b) $a=0.4959$ $b=0.9368$ (i) 2611 (ii) 5912 (iii) 3.237×10^{25} (c) 2954
8. (a) $\pi/128$ (b) $\sqrt{\pi}/12$ (c) (ii) $\ln \alpha$ and when $x=0$ $y=e^{-x(\ln x+c)}$
9. (a) $y = \tan(\tan x + c)$ (b) $y = e^{-x(\frac{1}{\alpha} + cx)}$ (c) $y = 1 + (-x^2 - c)^{-1}$
10. (a) $y = \frac{[(b-c)e^{ax} + (c-a)e^{bx} + (a-b)e^{cx}]}{[(a-b)(c-a)(c-b)]}$
 (b) $y = e^{ax}/(b-a)^2 - e^{bx}/(b-a)^2 + xe^{bx}/(b-a)$ (c) zero
11. Max at $x=1$; Min at $x=-1$ (a) zero (b) 1
12. (b) $\frac{1}{2} + \sum_{m=0}^{\infty} \frac{4}{(2m+1)^2 \pi^2} \cos((2m+1)\pi x)$

- II/1 : (a) (i) No (ii) $\phi = xy e^{z-x+x^2}$ (b) 69°
 2. $4\pi(R^2 - r^2 - \epsilon^2 - R)$ in both cases
3. (a) $\alpha = 1$. $x = 2/5$ $y = -7z/5$; $\alpha = -8/5$: $x = -5z$ $y = 5z/2$
 (b) (i) $S = I \cos \epsilon + \frac{A}{\epsilon} \sin \epsilon$ (ii) $T = I \cosh \epsilon + \frac{B}{\epsilon} \sinh \epsilon$
4. (a) Max $z = 1$ at $(1, 0)$; Min $z = -1$ at $(0, \pm 1)$
 (b) $x = (x_0 - Mct + My_0)/(1+M^2)$; $y = (Mx_0 + My_0 + c)/(1+M^2)$
 (c) $r = (3V/8\pi)^{1/3}$ $h = 2(3V/8\pi)^{1/3}$ (c) -
5. (a) see below (b) $\theta = \pi/6 + n\pi$ $n = 0, 1, 2, \dots$ or $\theta = n\pi$. $n = 0, 1, 2, \dots$ (c) -
6. (a) see below (b) $\Gamma \cdot \frac{1}{3}(1, 2, 2) = 20/3$ (c) 3 (d) (2, 4, 5) radius $\sqrt{7}$
7. (a) $F = U - TS$ (b) -
8. (a) exact, $f = 6x^2 + 5xy + 3y^2/2 - 9x - 4y + c$
 (b) not exact, $IF = x^{-1}r^2$ (c) exact, $f = -x/(x^2 + y^2)$
 (b) not exact, $IF = x^{-1}r^2$ (c) exact, $f = -x/(x^2 + y^2)$
9. (a) - (b) $\lambda = -1$ $(\sin 2\theta, -\cos 2\theta - 1, 0)$ ie $(\sin \theta, \cos \theta, 0)$
 $\lambda = 1$ $(\sin 2\theta, -\cos 2\theta + 1, 0)$ ie $(\cos \theta, \sin \theta, 0)$
 $\lambda = 2$ $(0, 0, 1)$
10. -
 11. (a) $5/2 \ln 2 - 1/2 \ln 3$ (b) $1/2$ (c) 1
12. $g' = g/t$ $g = kt$.

→ ∴ II/5(a) $\sqrt{2}, \pi/4, 4\sqrt{2}, \pi/8$

II/6(a) $|x-1|=4, (x-1)^2 + (y-2)^2 + (z-3)^2 = 16$

2006

II/1 : (a) $A=0$ (b)(i) no (ii) yes eg $B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ (c) $(I-D)^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$

2 : (a) (i) $\lambda = 0, \pm\sqrt{2}$ $\Sigma = \frac{1}{\sqrt{2}}(1, 0, -1), \frac{1}{\sqrt{2}}(1, \pm\sqrt{2}, 1)$ (ii) R is orthogonal
 $RMR^T = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sqrt{2} & 0 \\ 0 & 0 & -\sqrt{2} \end{pmatrix}$ M is not invertible ($\det M = \text{product of } \lambda's = 0$)
(b) $\lambda = 5, k_2(1, \cot\theta - 5\operatorname{cosec}\theta, k_1)$ with k_1, k_2 = arbitrary scalars

3. (a) (i) $-1/2 + i$ (ii) i (b) (i) $\pm\sqrt{3} - i, 2i$ (ii) $\pm(\sqrt{3} + i)$
(iii) $i e^{(\pi/2 + 2n\pi)} n=0, 1, 2, \dots$

4. (b) $(-2yz, 4xz^2, 0)$ (c) No, since $\nabla \cdot F \neq 0$ (d) $1/15$ in both cases

5. (a) $bc\dot{i} + ac\dot{j} + ab\dot{k}$ (b) $\int (bc\dot{i} + ac\dot{j} + ab\dot{k}) = abc$ (c) $abc/\sqrt{(b^2c^2 + a^2c^2 + a^2b^2)}$
(d) $\lambda + \mu + \nu = 1$ (e) $\lambda = b^2c^2/k, \mu = a^2c^2/k, \nu = a^2b^2/k$ where $k = b^2c^2 + a^2c^2 + a^2b^2$

(f) 90° .

6. (a) (ii) $(\frac{9}{16}, \frac{1}{2})$ (b) $(\frac{3\sqrt{3}R^3 \cos\theta_0 \sin^2\theta_0}{2})/2$

7. (a) (i) 0.05094 (ii) 0.0194 (iii) 1.05×10^{-5} (b) (i) $5/12$ (ii) $5/9$

8. (a) Max at $(0, 1)$ $f = 2$ Min at $(0, -1)$ $f = -2$ (b) $(1, -1/2, 1/2)$

9. (a) $f(a+x) = f(a) + xf'(a) + \frac{x^2}{2!} f''(a)$

(b) (i) $x - x^2/2 + x^3/3 - x^4/4$ (ii) $\pi/4 + x/2 - x^2/4 + x^3/12 + \dots$ (iii) $1 - x/2 + x^2/12 + \dots$ Yuk

10. (a) $y = \frac{1}{2}x^3(\cos x - \cos 2x)$ (b) $y = \frac{1}{8}\cos x - \frac{1}{8}\cos 2x + \frac{1}{8}\sin 2x + \frac{1}{8}$

(c) $y = (\frac{1}{2}x^2 + \frac{1}{6}x^3)e^x$

11. (b) $\sum_{n=0}^{\infty} [8/\pi(4-(2n+1)^2)] \cos(2n+1)x$

12. (b) $f = 0$ everywhere

II/2 : (a) $(1+x^2)^{-1}$ (b) $x^2/18$ (c) $(32/15)/3^{1/4}$

2. (b) $4(\frac{a}{2}, \frac{b}{2})(b_i a_j - a_i b_j)$ (c) Show that $(CBA)_{ij} = -\delta_{ij}$ & $S''' = -S$

3. (a) F_1 and F_3 (b) $\pi^4/16$

4. (a) $\bar{F} = \underline{\operatorname{grad}} \underline{F} / (\partial \underline{F} / \partial \underline{x})$ (b) $\int (2x(1+y), x^2+2y^2, y^2)/y^2 dx dy$ (c) 2

5. (a) $\frac{1}{2}\pi a^3$ (b) $\pi a^3/(1-\cos\theta_0)^2 (1-2\cos\theta_0) \sin^3\theta_0 d\theta$ (slice parallel to xy plane) (c) $5\pi a^3/2$

6. (a) $\frac{2}{x} \cos x^4 - \frac{1}{x} \cos x^3$ (b) bounds are $1-h/2$ and 1

7. (a) $x^2y^2 + 4x - 6y = c$ (b) $(y^2 + x - 1)e^x = c$ (c) $x \cos x + xy^2 = c$

8. (a) (ii) all $\frac{1}{4^4}$: $(1, 12, 54, 108, 81)$ (b) (i) $3 \cdot 5 (ii) 15^{1/6} (iii) 2^{11/12}$

(c) (i) $x = 1/4$ (ii) $2 \cdot 58/3$

9. (b) $z = e^{a(x+c)} - e^{b(x+c)}$ (c) $y = \frac{1}{x+c} + a \cdot z = (x+c)e^{ax+c}$

(d) $z'' - (a+b)z' + abz = 0$ (can also get expressions for $c-x$ for z if you treat log integral differently)

10. (b) $\lim_{n \rightarrow \infty} \omega_n = 0$

11. Yuk

12. Max is at $(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$; Min at $-(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$.

- I/A 1. $x=1, i\sqrt{2}, -2$ 2. (a) $-5\sin 5x$ (b) $2xe^x + x^2e^x$ 3(a) $(3+2x)^{1/2} + c$
 3(b) $\frac{1}{2}x^2 \ln x - \frac{1}{4}x^2 + c$ 4. radius 2 centre $(0, -3)$ 5(a) 10201 (b) $3/2$
 6. 6 7. $\pi/2$ or $\pi/4$ 8. 30° 9. 5 10. - 11. $k=2e$ $x = \sqrt[3]{2}$ $y = e$

B1. -

2. (a) $Re = \pm \frac{1}{\sqrt{2}}$ $Im = \pm \frac{1}{\sqrt{2}}$; $Re = 1$ $Im = \pi$; $Re = \cosh 2$ $Im = \sinh 2$; $Re = 0$, $Im = -1$

. (b) $\sin 3x = 3 \sin x - 4 \sin^3 x$ $x = 0, \pi, \pi/6, 5\pi/6, 7\pi/6, 11\pi/6$
 (c) $-b \pm \sqrt{b^2 - 1}$ roots trace out unit circle, centred on origin

3. (a)(i) $x - \frac{1}{3}x^3$ (ii) $+x \ln a + \frac{x^2}{2} \ln^2 a$ (iii) $-x^2 - x^4/6$ (b) 3rd of 6 terms

4. (a) - (b) 0.31, 0.0362.

5. (a) (i) exact: $(1+ey)\sin x = c$ (ii) Not exact, $\mu = \frac{1}{y}$: $\frac{x^3}{3} + x \ln y = c$
 (b) $y = (1+x)^5/2 + (1+x)^3$

6. $(0,0)$, $(\pi/2, \pi/2)$, (π, π) , $(0, \pi)$, $(\pi, 0)$
 \min saddle \min \max \max

7. (i) $\frac{1}{4}(e^4 - 1)$ (ii) Mass = $\pi k/4$ Area = $4\pi(1 - 1/\sqrt{2})$

8. (a)(i) False (ii) True, $a+b$ (iii) True, ab (b)(i) 0, 1 (ii) 0, 1

(c) $-1, 0, 3$; $(1, 0, -1)^T$.

9. (a) $\sum \frac{1}{n} \ln n$ diverges; $\sum \ln \frac{1}{n}$ converges: $\ln^{-1}(N+1) < R_N < \ln^{-1}(N)$

(b) converges; diverges.

10. I'd avoid ever trying...

- II/A 1. $(-2, 0, 2)$ 2. (a) 250 (b) $i\sqrt{2}$ 3. (b) $(-1, -1)$ 4. (a) $-(6x, 6y, 2z)$
 4(b) -14 (c) $\pm \infty$ 5. (a) 5 (b) $\begin{pmatrix} 2 \\ 1 \\ 6 \\ 3 \end{pmatrix}$ 6. $M\omega = \sqrt{8}$ Arg = $5\pi/12$
 7. $x - x^2/2$ 8. $t = \frac{1}{k}(\ln x_0)$ 9. (a) $4/7$ (b) $6/7$.

B1. $r = 2\cos\theta$; $r = 2/\cos\theta$

2. -

3. (a) $A = (-\rho; \rho^{n+1}; \rho^{n-m})$ (b) $B = \lambda; e^{\lambda(y-x)}; \lambda e^{\lambda(y-x)}; 1/\lambda^2$

4. (a) $y = (A+Bx)e^{-2x} + \frac{1}{16}e^{2x}$ (b) $y = -\frac{5}{4}e^{-3x}\sin 4x + \sin 5x$

5. (a) $(\partial u / \partial y)_x = (\partial v / \partial x)_y$ (b) $(\partial v / \partial x)_y$; both sides equal to $-2T/3V$.

6. (a) Ω (b) $\varphi = -(x^2y + xz + y^2 + c)$ (c) 3 (d) $4\pi b^2 a^3/15$

7. (a)(i) True (ii) False (iii) True (b) $C^{16} = C$ (c) $\mu \neq 0$ $M^{-1} = \frac{1}{2\mu} \begin{pmatrix} 1 & \mu & -1 \\ -1 & \mu & 1 \\ 1 & -1 & \mu \end{pmatrix}$

7. (d)(i) $x=1$ $y=1-\mu$ $z=1$ (ii) $y=1$ $x+z=2$

8. $f(x) = \frac{1}{2} + \sum_{n=1}^{\infty} \left(\frac{2}{n^2\pi^2} ((-1)^n - 1) \cos n\pi x - \frac{2}{n\pi} (-1)^n \sin n\pi x \right)$

9. stationary pt. is $(1, 1)$; contour parallel to constraint path at stationary point.

10. $\frac{dT}{dt} = aVt$; $u = e^{-\pi^2 t} \sin \pi y$; $u = e^{-\pi^2 t} \sin \pi y + \frac{1}{10} e^{-9\pi^2 t} \sin 3\pi y$

2008 (Question numbering for online version)

- I/A 1. $\theta = 75^\circ \text{ or } 345^\circ$ 2. $2x\cos x^2$; $\frac{e^{2x}}{x} (2 - 4x)$ 3. $(x^2+1)^{1/2} + C$, $(x-1)e^{x+C}$
 4. $x = \pm 1, \pm \sqrt{2}$ 5. $\sqrt{10}$ 6. $y = 3-x$ 7. $x < 1 \text{ or } x > 2$ 9. $\sin \theta + \theta$
 10. (i) $(N+1)^3$ (ii) $\frac{1}{3}((N+1)^3 - (N+1))$
- BII. $\underline{n} = \frac{1}{\sqrt{3}}(\underline{i} + \underline{j} + \underline{k})$; $\underline{e}_1 = \frac{1}{\sqrt{6}}(1, 1, -2)$; $\underline{e}_2 = \frac{1}{\sqrt{2}}(1, -1, 0)$ ratio = $\sqrt{3}$.
 coords $\frac{1}{\sqrt{6}}(x+y-2z)$, $\frac{1}{\sqrt{2}}(x-y)$

12. (b) (i) $\frac{1-x^2-y^2}{(1+x)^2+y^2}$, $\frac{-2y}{(1+x)^2+y^2}$ (ii) $e^{-y}\cos x$, $e^{-y}\sin x$ (iii) Re: $x \sin x \cosh y$,
 IM: $-y \cos x \sinh y$, $+x \cos x \sinh y$
 (c) (i) $e^{i\theta}$ where $\theta = \pi/3, 2\pi/3, 4\pi/3, 5\pi/3$ (ii) $i(\cosh 2 \pm i \sinh 2)$
13. (a) $2(x + x^{3/3} + x^9/5 + \dots)$ (b) $(-19/216)x^4$ (c) $a = 2/3$, $b = 1/6$
14. (a) $1/6(17^{3/2}-1)$ (b) volume is $(p-q)\pi r^3$; area is $2(p-q)\pi r^2$.
15. (b) $P(x < z < \beta) = \int_x^\beta \sum_{n=-\infty}^{\infty} p_n f(z-n) dz$; density = $\sum_{n=-\infty}^{\infty} p_n f(z-n)$
16. (a) (i) $y = (-x)^3$ (ii) $y = 1/2 \ln(1+x^2)$ (iii) $y = x \sin(\ln x + c)$
 (b) (i) $y = 2 \sin x + 2e^{-x} \cos x$ (ii) $y = -bx$??
17. (a) (i) $(0, 1/4), (1/4, -1/4), (-1/4, 0)$ (ii) $(1/3, -1/3)$, ~~minimum~~
 (b) saddle at $(0, 0)$; minimum at $(-2/3, 0)$
18. (a) (i) -4 (ii) $\frac{1}{4} \begin{pmatrix} 1 & -2 & 3 \\ -2 & 0 & 2 \\ 3 & 2 & -3 \end{pmatrix}$ (iii) $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ (b) - (c) $\mu = 5$.

19. (a) (i) all $n > 0$ (ii) $n=1, 2, 3$ (iii) $n=2, 3$ (b) $\max(A, B)$ (c) nothing; converges.

20. $V_x > V_t$.

- II/A 1. $\pi/3$ 2. (i) 1 (ii) 0 3. $x = \pm y$ 4. (i) 2π (ii) $5\pi^2$ 5. $(0, 1)$ 6. $\pm 3/2$
 7. $e^{-ex^2/2}$ 8. (i) $\alpha = 1$ (ii) $u = x^2 - x + \epsilon$ 9. (i) $\pi/4$ (ii) $8\pi dr$

B10. -

11B. (a) (i) $\frac{L!(N-S)!}{(L-S)! N!}$ (ii) $\frac{S!}{n!(S-n)!} \cdot \frac{L!(N-L)!(N-S)!}{(L-n)!(N-L-S+n)! N!}$ (b) $\frac{L}{N}$ (c) $\frac{K!}{k!(K-k)!} f^k (1-f)^{K-k}$

(d) (i) $1/4$ (ii) $4/100$ (iii) $0.1, 0.735$

12B. $I(a, b) = \int_a^b e^{-x^2} dx$ (checked using online definite integrator!) $J(a, b) = I(b, a)$

13B. (i) $y = -16e^{3x} + 9e^{4x} + 12x + 7$ (ii) $y = 5e^{-x} - 2e^{-2x} + \sin x - 3\cos x$

(iii) $y = 4x^2 e^{-x}$

14B. (a) $v = 4$ $A = -2$ (b) -

15B. (a) (i) ϕ (ii) ϕ not (b) (i) $F_1 = -1$ $F_2 = -50$ (ii) $F_1 = -1$ $F_2 = -323/7$

16B. (a) (i) A^2 (ii) A (iv) A^2 (b) $\mu = 4$ $\lambda = 1 : \frac{1}{\sqrt{5}}(-2, 0, 1)$ $\lambda = -4 : \frac{1}{\sqrt{5}}(2, -5, 4)$

$\lambda = 5 : 1/3(1, 2, 2)$ (ii) $|\mu| < 1$

(iii) $\sum (-2(-1)^n \sin nx)/n$

17B. (a) $2L$ (b) - (c) $\sum (-2(-1)^n \sin nx)/n$

18B. (a) $4\pi a^2 b$ (b) $4\pi a^2 b \frac{K}{3}$

19. (a) $a_n = \frac{2L}{n\pi c} \int_0^L v(x) \sin \frac{n\pi x}{L} dx$ frequencies $n=1$ and $n=3$
 ratio of amplitudes $3:1$.

(b) $f(x) = A \cos kx + B \sin kx$; $v(x, y) = -\pi \sum_a \cos x e^{-ly}$

2009

Paper I

- A1. $-1 \pm \sqrt{3}$. 2. $x = 0$ or 9. 3. (a) $3/x$; (b) $\cos 2x - 2x \sin 2x$.
4. (a) $\frac{1}{4}(\ln|x-4| - \ln|x|) + c$; (b) $\frac{1}{2}e^{x^2} + c$. 5. $x = \frac{\pi}{2}$ or $\frac{2\pi}{3}$. 6. $x = 25$. 7. (a) 15352 (b) 4/5. 8. - 9. $16/\sqrt{5}$. 10 (b) -2.
- B11. (a) (i) $\frac{-11x^3 - 6x^2}{(3x-2)^5}$; (ii) $a/(a^2 - x^2)$; (c) (i) $\pi(h-x)^2 x \tan^2 \alpha$; (ii) $\frac{3}{4} \cot \alpha$.
12. (a) (i) $\binom{n}{k} p^k (1-p)^{n-k}$; (ii) $np, np(1-p)$; (b) (i) $\binom{n-1}{c-1} p^c (1-p)^{n-c}$.
13. (a) $3 \pm i, \begin{pmatrix} 2 \\ -i-1 \end{pmatrix}, \begin{pmatrix} 2 \\ i-1 \end{pmatrix}, B = \begin{pmatrix} 2 & 2 \\ -i-1 & i-1 \end{pmatrix}$; (b) $-4, 1, 6, \begin{pmatrix} 4 \\ -5 \\ 3 \end{pmatrix}, \begin{pmatrix} 3 \\ 0 \\ -4 \end{pmatrix}, \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix}$.
14. (b) $e^{i\theta}$, where $\theta = \frac{\pi}{8} + \frac{n\pi}{2}, n = 0, 1, \dots$; If you regard the square root of i as only having one value (rather than being +/- that value) then you will get $\theta = \frac{\pi}{8} + n\pi$.
15. -
16. (b) 24; (d) $\pi, \sqrt{\pi}$.
17. (a) (i) $\sin\left(\frac{y}{x}\right) = \frac{2}{\pi x}$; (ii) $x = \tan\frac{(x+y)}{2}$ (b) $y = \operatorname{cosec}^2 x$.
18. (a) $(\frac{\partial u}{\partial s})_t + (\frac{\partial u}{\partial t})_s (\frac{\partial t}{\partial s})_v$; (b) $\mu = Ax^{-3}, f = -x^{-2} \sin y + c$.
19. (c) $\sum_{n=1}^{\infty} \sin \frac{n\pi x}{l} (C_n \cos \frac{n\pi ct}{L} + D_n \sin \frac{n\pi ct}{L})$.
20. (a) (i) No; (ii) Yes; (iii) No; (b) (i) 3; (ii) 4; (iii) 1/90.

Paper II

- A1. - 2. Both zero 3. $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ 4. $\alpha = \pm 1$ 5. Max at $x = 0$, Min at $x = 2$ 6. -
7. $y = x - 1$ 8. Ignoring their (apparently daft) suggestion and finding the Taylor Series directly gives $\ln 2 + \frac{x}{2} + \frac{x^2}{8}$. 9. 0 to $\frac{\pi}{4}, \frac{3\pi}{4}$ to $\frac{5\pi}{4}, \frac{7\pi}{4}$ to 2π .
- B11. (a) $(1, 2, 4) + \lambda(1, 2, -2)$ or $x = 1 + \lambda, y = 2 + 2\lambda, z = 4 - 2\lambda$;
 (b) $(1, -1, 0) + z(2, 3, 1)$; (c) $\sqrt{\frac{35}{49}}$.
12. (ii) $y = Ae^{\frac{t(-\lambda+\sqrt{\lambda^2-4k})}{2}} + Be^{\frac{t(-\lambda-\sqrt{\lambda^2-4k})}{2}}$, oscillatory if $\lambda^2 < 4k$; (ii) $\omega = \sqrt{\frac{4k-\lambda^2}{2}}, t = \frac{2}{\lambda} \ln \frac{4}{3}$; (iii) $\frac{1}{(1+\omega^2)^2} \sqrt{16 + \left(\frac{2-2\omega^2}{\omega}\right)^2}$???
13. (a) (1,0) saddle, (0,1) min, (0,-1) max, (2,1) max, (2,-1) min; (b) (i) $\frac{1}{\sqrt{2}}(1, 0, -1)$;
 (ii) $\frac{u}{\sqrt{2}}(-1, 0, 1)$; (iii) (0,1).
14. (a) (i) -4; (ii) $\frac{1}{4} \begin{pmatrix} 3 & -5 & 1 \\ 1 & 1 & -1 \\ 1 & -3 & -1 \end{pmatrix}$; (b) 5; (c) $b = a$ or $c = a$ or $c = b$.
15. (b) $\frac{\pi^4}{90}$; (c) $A_0 = 1; A_n = 2(-1)^n \cos n\pi x / (n^2\pi^2 + 1)$.
16. (c) $\underline{E} = \frac{\underline{r}}{r^3}; \nabla \times \underline{E} = \underline{0}$.
17. (c) (i) $P(A \cap B \cap C)$; (ii) $P(A)$; (iii) $P(B)$; (iv) $P(A \cup B)$; (v) $P(B|A)$; (d) $N = 3744$.
18. (a) $e^2 - 3$; (b) $\frac{1}{6}$; (c) $\pi - \frac{2}{3}$.
19. (a) zero, $\frac{\pi}{4}$; (b) zero.

